

Targeting Resistance in EGFRm NSCLC with HER3-Directed ADCs in the Community Setting

This transcript has been edited for style and clarity and includes all slides from the presentation.




Provided by

Targeting Resistance in EGFRm NSCLC with HER3-Directed ADCs in the Community Setting

Helena Yu, MD






► **Dr. Yu:** Hello and welcome to this educational activity. My name is Helena Yu. I am the research director of the Thoracic Oncology Service as well as an associate attending and medical oncologist at Memorial Sloan Kettering Cancer Center in New York. Today we will be discussing HER3-directed antibody-drug conjugate therapies for the treatment of EGFR-mutant non-small cell lung cancer. So, let's get started.



► EGFR-mutant lung cancer represents 15 to 20% of all non-small cell lung cancer and the different driver mutations we see in non-small cell lung cancer you can see on the left. A lot has happened in regards to drug development in the last 22 years. The EGFR mutation was discovered in 2004 and since that time various generations of EGFR inhibitors have been developed. We've begun to understand mechanisms of resistance to these EGFR inhibitors and more recently we've seen the advent of antibody-drug conjugate therapies for EGFR-mutant lung cancer.

1L Treatments for EGFR-Mutant Advanced Lung Cancer



*Currently not an approved first-line combination. EGFR, epidermal growth factor receptor; HR, hazard ratio; PFS, progression-free survival. Soria J-C, et al; for the FLAURA Investigators. *N Engl J Med.* 2018;378:113-125. Ramalingam SS, et al. ESMO 2019. Abstract 567.

► In regards to first-line treatment for EGFR-mutant lung cancer, the standard of care would be osimertinib, which is a third-generation EGFR tyrosine kinase inhibitor. It was compared to earlier generation EGFR TKIs in the FLAURA study and was shown to have better progression-free survival as well as overall survival compared to earlier generation TKIs.


More recently, in the last year, there have been EGFR

TKI-based combinations that have been assessed for first-line treatment. So one such regimen is the FLAURA2 regimen which is adding in platinum-based doublet chemotherapy to osimertinib as first-line treatment and that also demonstrated a progression-free survival benefit compared to osimertinib alone. And then there's the combination of amivantamab and lazertinib. Amivantamab is an EGFR-

MET bispecific antibody, and lazertinib is a different third-generation EGFR TKI and that combination was also shown to be superior to osimertinib in regards to progression-free survival. We don't yet have the overall survival data for these combinations, but both are likely to be approved options that we could consider for our patients.

Mechanisms of Resistance to Osimertinib

- Mechanisms of resistance to first-line osimertinib are diverse and no one mechanism is dominant **so upfront combinations to prevent resistance not appropriate without a biomarker**
- With development of better EGFR inhibitors, there is more off target resistance seen
- High incidence of lineage plasticity including both small cell and squamous transformation
- Frequent acquired gene alterations such as gene fusions which are rare de novo
- There will be a role for non-biomarker selected therapies that focus on enhanced EGFR on-target inhibition or address general tumor biology




EGFR, epidermal growth factor receptor.
Jordan EJ, et al. *Cancer Discov.* 2017;7(6):596-609.

▶ So although all patients respond to osimertinib, everyone subsequently develops resistance to therapy. We biopsy patients at the time of resistance to understand why the tumor has become resistant to targeted therapy. When we biopsy patients there can be different subsets of resistance. One such subset is on-target resistance, which means


acquired alterations in EGFR like EGFR C797S. Resistance mutations can be off-target, which means acquired alterations or amplification in different bypass signaling pathways. And then, interestingly, we can see something called histologic transformation, which is when an adenocarcinoma becomes a squamous cell lung cancer or a small cell lung cancer and

this change in histology drives resistance to targeted therapy. About half the time we don't see a genomic alteration or histology change that drives resistance. And so, with such diverse resistance mechanisms, a targeted therapy that would be efficacious in all types of resistance is something that would be very valuable.



▶ HER3 is part of the HER protein kinase family, which includes other driver mutations like HER2 as well as EGFR. The HER family members heterodimerize with HER3 so there isn't signaling from the HER3 receptor alone, it needs to heterodimerize or pair with other receptors, and that heterodimerization leads to down-stream signaling that leads to self-proliferation, cancer cell survival.

HER3 expression also can mediate resistance to targeted therapy where we see up-regulation of HER3 in the acquired resistance setting.




▶ In terms of HER3 expression in non-small cell lung cancer, first we don't tend to see HER3 mutations or genomic alterations within non-small cell lung cancer, but we do tend to see HER3 protein expression and that's measured by immunohistochemistry. So, the way that we use IHC is we calculate an H-score which tells us the intensity of staining as well as the percent of cells that are positive and you can see some IHC staining for

HER3 in the top right. When we look at HER3 expression within non-small cell lung cancer the degree of HER3 expression is related to both progression or recurrent disease as well as metastatic disease and so it is a negative prognostic biomarker within non-small cell lung cancer. However, because we aren't currently using HER3 expression in a clinical manner, HER3 testing by IHC is not currently recommended as standard of care.

So, when we think about the relevance of these novel HER3-directed therapies, I think that it is very exciting to think about how these different antibody-drug conjugates, including HER3-directed therapies, might be integrated into our clinical practice. Post osimertinib first-line treatment, we don't have any targeted therapies that are currently approved and so to have additional options besides standard cytotoxic chemotherapy would be really helpful.

Patritumab Deruxtecan

- HER3-DXd is an ADC composed of 3 parts¹⁻⁴:
 - A fully human anti-HER3 IgG1 mAb (patritumab)
 - A topoisomerase I inhibitor payload (an exatecan derivative, DXd)
 - A tetrapeptide-based cleavable linker that covalently bonds the other 2 components



7 Key Attributes of HER3-DXd

- Payload mechanism of action: topoisomerase I inhibitor^{1-4,a}
- High potency of payload^{1-4,a}
- High drug-to-antibody ratio ≈ 8 ^{1,2,a}
- Payload with short systemic half-life^{2,3,a,b}
- Stable linker-payload^{2-4,a}
- Tumor-selective cleavable linker^{1-5,a}
- Bystander antitumor effect^{2,6,a}

AXIS
Medical Education

^aThe clinical relevance of these features is under investigation. ^bBased on animal data.
ADC, antibody-drug conjugate; HER3-DXd, patritumab deruxtecan; IgG1, immunoglobulin G1; mAb, monoclonal antibody.
1. Hashimoto Y, et al. *Clin Cancer Res.* 2019;25:7151-7161. 2. Nakada T, et al. *Chem Pharm Bull (Tokyo).* 2019;67(3):173-185. 3. Ogitani Y, et al. *Clin Cancer Res.* 2016;22(20):5097-5108. 4. Koganemaru S, et al. *Mol Cancer Ther.* 2019;18:2043-2050. 5. Haratani K, et al. *J Clin Invest.* 2020;130(1):374-388. 6. Ogitani Y, et al. *Cancer Sci.* 2016;107(7):1039-1046.

► The first-in-class medication is one called patritumab deruxtecan. The other name for it is HER3-DXd, which is an antibody-drug conjugate that is composed of three parts. HER3-DXd has a fully human anti-HER3 monoclonal antibody, patritumab, and then ADCs always have a

chemotherapy payload and in this case it's a topoisomerase I inhibitor payload, so an exatecan derivative. The payload and the antibody are combined, bound together by a cleavable linker. And so when we talk about ADCs, one terminology that we might hear of is something called the


drug-to-antibody ratio and that really describes how many molecules of a chemotherapy payload are attached to a single antibody. And so, the higher the drug-to-antibody ratio, theoretically the more potent the ADC might be.

HERTHENA-Lung 01 Phase 2

HER3-DXd IV Q3W

Patient population

- Advanced *EGFR*-mutated NSCLC
- Progression on most recent systemic therapy
- Prior *EGFR* TKI and prior platinum-based chemotherapy (amended protocol required prior osimertinib)
- Inactive or previously treated asymptomatic brain metastases allowed
- Pretreatment tumor tissue required^a



^aProvided as either: Pretreatment tumor biopsy from at least 1 lesion not previously irradiated and amenable to core biopsy; or archival tumor tissue collected from a biopsy performed within 3 months prior to signing of the tissue consent and since progression while on or after treatment with the most recent cancer therapy regimen

Baseline characteristics		HER3-DXd 5.6 mg/kg (N=225)
Age, median (range), years		64 (37-82)
Female, n (%)		132 (59)
Asian, n (%)		105 (47)
Time since initial NSCLC diagnosis, median (range), months		41.0 (9.1-224.7)
Sum of target lesion diameters at baseline (BICR), median (range), mm		68 (11-248)
History of CNS metastasis, n (%)		115 (51)
Brain metastasis at baseline (BICR), n (%)		72 (32)
<i>EGFR</i> -activating mutations, n (%)	Ex19del	142 (63)
	L858R	82 (36)
No. of prior lines of systemic therapy (locally advanced/metastatic)	Median (range)	3 (1-11)
	2 prior lines, n (%)	58 (26)
	>2 prior lines, n (%)	165 (73)
Prior cancer regimens, n (%)	Prior <i>EGFR</i> TKI therapy	225 (100)
	Prior third-generation <i>EGFR</i> TKI	209 (93)
	Prior platinum-based chemotherapy	225 (100)
	Prior immunotherapy	90 (40)

BICR, blinded independent central review; CNS, central nervous system; cORR, confirmed objective response rate; DOR, duration of response; *EGFR*, epidermal growth factor receptor; HER3-DXd, patritumab deruxtecan; IV, intravenous; NSCLC, non-small cell lung cancer; Q3W, once every 3 weeks; TKI, tyrosine kinase inhibitor.
Yu H. et al. WJCL 2023. Abstract OA05.03.


► The first study to look at HER3-DXd was a first-in-human phase 1 study that helped assess the ideal dose for the phase 2 study. And here you can see the schema for the HERTHENA-Lung01 study, which is the registrational phase 2 study. The phase 1 study really demonstrated that HER3-DXd can be given safely, and it was focused on patients with *EGFR*-mutant lung cancer. Similarly, HERTHENA-Lung01 assessed patients with *EGFR*-mutant lung cancer that had progressed on the most recent systemic therapy. They needed to have been previously treated with both an *EGFR* targeted therapy as well as prior platinum-based chemotherapy. Patients with asymptomatic treated or

untreated brain metastases were allowed to enroll and, while HER3 wasn't assessed prospectively, pretreatment tumor tissue was required for retrospective assessment of HER3 expression.


So the study initially consisted of 2 cohorts: one with a fixed dose of HER3-DXd of 5.6 mg/kg given intravenously every 3 weeks, as well as an up-titration cohort where increasing doses of HER3-DXd were assessed.

After enrollment of a small number of patients based on benefit-risk assessment of the phase 1 data of these different dosing schedules, the fixed dose of 5.6 mg/kg was chosen to complete accrual. So, a total of 225 patients were enrolled at the fixed dose of

5.6 mg/kg and the primary endpoint of the study was confirmed overall response rate by independent central radiology review. You can see the table on the bottom right shows the baseline characteristics of patients on study and they really were consistent with what we see for *EGFR*-mutant lung cancer with an enrichment in female patients as well as Asian race. Notably, more than half of patients had a history of CNS metastases, which is quite common in *EGFR*-mutant lung cancer and the study focused on the 2 common sensitizing mutations, exon 19 deletions as well as L858R, and all patients had received prior *EGFR* TKI as well as platinum-based chemotherapy.




► And here you can see the waterfall plot that shows the percentage of disease shrinkage of target lesions on her HER3-DXd. And you can see that the majority of patients did have tumor shrinkage on therapy. The confirmed overall response rate with maximal follow-up was 29.8%. And then on the waterfall plot you can see that patients had different mechanisms of resistance to previous EGFR TKI including EGFR independent, dependent, and unidentified resistance mechanisms and you can see that HER3-DXd was effective in all of those different scenarios.



► Importantly, the study authors wanted to assess intercranial efficacy. As I mentioned, this patient population has more than a 50% cumulative incidence of brain metastases. And so, they identified 30 patients that were treated on study that had brain metastases at baseline but had not received prior radiotherapy and so these were measurable CNS lesions. And when looking at intercranial response rate, that was 33.3%, so very similar to the systemic or overall response rate, really demonstrating that this drug has CNS activity. And you can see from the CT images on the right, where you can see complete resolution of a lesion while on therapy.

HERTHENA-Lung 01 Phase 2

Association of Baseline Tumor HER3 Membrane H-Score With Confirmed BOR by BICR Following Treatment With HER3-DXd 5.6 mg/kg (N=225)^a



BICR, blinded independent central review; BOR, best overall response; CR, complete response; HER3, human epidermal growth factor receptor 3; HER3-DXd, patritumab deruxtecan; NE, not established; PD, progressive disease; PR, partial response; SD, stable disease. Yu H, et al. WCLC 2023. Abstract OA05.03.

► So, one important thing to understand is to see whether response to HER3-DXd is associated with degree of HER3 expression. So one thing that's important to note is that this study focused on EGFR-mutant lung cancer and the majority of EGFR mutant lung cancers do express some degree of HER3. So there were very few HER3-negative patients that were enrolled. But when you separated out

patients that had a response to therapy, stable disease, or progressive disease, you can see there is complete overlap of HER3 expression between those three groups and so the degree of HER3 expression really does not predict response to HER3-DXd.

When we think about toxicity the important thing to think about with antibody-drug conjugates is that they are a hybrid between targeted


therapy and chemotherapy. So they can have chemotherapy-like side effects and that would include cytopenias including thrombocytopenia and neutropenia and anemia, as well as sometimes nausea, vomiting, and alopecia. And then sometimes there can be toxicities that are associated with the antibody, which might be side effects similar to what we see with targeted therapy.

HERTHENA-Lung 01 Phase 2

Safety summary	HER3-DXd 5.6 mg/kg (N=225)
Any TEAE, n (%)	224 (99.6)
Associated with treatment discontinuation	16 (7.1)
Associated with treatment dose reduction	48 (21.3)
Associated with treatment dose interruption	91 (40.4)
Associated with death	24 (10.7)
Grade ≥3 TEAE, n (%)	146 (64.9)
Treatment-related TEAE, n (%)	215 (95.6)
Associated with death	4 (1.8)
Grade ≥3	102 (45.3)
Serious TEAE	34 (15.1)
Adjudicated interstitial lung disease, n (%) [All were adjudicated as treatment-related]	12 (5.3)
Grade 1	1 (0.4)
Grade 2	8 (3.6)
Grade 3	2 (0.9)
Grade 4	0
Grade 5	1 (0.4)

Median treatment duration: 5.5 (range, 0.7-18.2) months.

Most Common Grade ≥3 TEAEs Occurring in ≥3% of Patients (N=225)



Any hematologic toxicities typically occurred early in treatment, were transient, and were not associated with clinical sequelae




HER3-DXd, patritumab deruxtecan; TEAE, treatment-emergent adverse event.
Yu H, et al. WCLC 2023. Abstract OA05.03.

Importantly, when assessing the safety of a clinical trial and a novel therapy I look at the rate of treatment-emergent adverse events that were associated with treatment discontinuation and that was relatively low on this study at 7.1%. You can see that there was cytopenias that were seen. Those were mostly front-loaded in the early part of the study and very few were related or associated with clinical sequelae like clinical bleeding or febrile neutropenia. One important class effect of these antibody-drug conjugates is something called interstitial lung disease or pneumonitis which we know can occur with chemotherapy, can occur with targeted therapy, but does also occur with these deruxtecan backbone antibody-drug conjugates. This was looked at by an independent adjudication committee and the rate of ILD or pneumonitis on study was 5.3%.

HERTHENA-Lung02: Ongoing Phase III Study of Patritumab Deruxtecan in EGFR-Mutated NSCLC

Multicenter, randomized, open-label phase III study

Stratified by prior third-generation EGFR TKI (osimertinib vs other; 1L vs 2L); region (Asia vs RoW), brain metastases (yes vs no)



- Primary endpoint: PFS by BICR (RECIST v1.1)
- Secondary endpoints: PFS by investigator, OS, ORR, DoR, DCR, TTR, safety



1L/2L, first-line/second-line; BICR, blinded independent central review; CT, chemotherapy; DCR, disease control rate; DoR, duration of response; EGFR, epidermal growth factor receptor; HER3, human epidermal growth factor receptor 3; HER3-DXd, patritumab deruxtecan; IV, intravenous; NSCLC, non-small cell lung cancer; ORR, objective response rate; OS, overall survival; PFS, progression-free survival; Q3W, once every 3 weeks; RECIST, Response Evaluation Criteria in Solid Tumours; TKI, tyrosine kinase inhibitor; TTR, time to relapse.
Mok TSK, et al. Future Oncol. 2024;20(15):969-908.
ClinicalTrials.gov identifier: NCT05338370.

So, the registrational phase 2 study is complete with results that had been already presented. What is ongoing is this confirmatory phase 3 study that is focusing on the same patient population, so patients with metastatic EGFR-mutant lung cancer that have received prior EGFR TKI. The notable difference is that this is prior to patients receiving platinum-based chemotherapy. So they were allowed to receive TKI and then they were randomized 1:1 to receiving HER3-DXd versus platinum-based chemotherapy, which would be the standard of care off-study. And the primary endpoint of this study is progression-free survival by independent central radiology review, and we haven't yet seen results for the study. It's ongoing.

Patritumab Deruxtecan Regulatory Status

- BLA seeking accelerated approval granted priority review by FDA on 12/22/2023¹
- FDA issued a CRL on 6/25/2024²
 - The CRL results from findings **pertaining to an inspection of a third-party manufacturing facility**
 - **The CRL did not identify any issues with the efficacy or safety data submitted**
- The companies developing patritumab deruxtecan are working with the third-party manufacturer to address and resolve the findings to continue the approval process²

► So, when we think about or when we note the status of patritumab deruxtecan it did have a breakthrough designation and was seeking accelerated approval and the approval of this drug was expected but we note that in June of 2024 this year, the FDA issued something called a complete response letter. And this is important to note that this CRL, this complete response letter, did not identify any issues with the efficacy or safety data submitted, but the drug approval was delayed due to an inspection of the third-party manufacturing facility. And so, we know that remedy of this situation is currently ongoing, and we do expect approval of HER3-DXd in the future.


► So, HER3-DXd is not the only HER3-directed antibody-drug conjugate that is currently in development. There is another agent called BL-B01D1 which is a bispecific EGFR-HER3 antibody-drug conjugate. And the initial data was from the phase 1 study that was done in China and that looked at various different solid tumors including lung cancer that was both EGFR-mutant as well as EGFR wild-type. But when focusing on the EGFR-mutant non-small cell lung cancer they treated 40 patients and the response rate was 67.5%. So really a meaningful clinical efficacy in this patient population. And based on this phase 1 completed in China, there is an ongoing global study assessing BL-B01D1 in different solid tumors including EGFR-mutant lung cancer.



BLA, biological license application; CRL, complete response letter; FDA, Food and Drug Administration.
 1. Daiichi Sankyo, Inc. https://www.daiichisankyo.com/files/news/pressrelease/pdf/202112/20211223_E1.pdf
 2. Daiichi Sankyo, Inc. https://www.daiichisankyo.com/files/news/pressrelease/pdf/202406/20240620_E.pdf

BL-B01D1: Response Rates in EGFR-Mutated NSCLC

- All patients in the current analysis received Q3W dose regimens



	EGFR-Mutated NSCLC		EGFR Wild-Type NSCLC	
	All (n = 40)	Treated/No CNS Mets (n = 13)	All (n = 62)	2L Post PBC (n = 26)
Prior CT lines, %				
• 0	25	8	0	0
• 1	50	46	42	100
• 2+	25	46	56	0
ORR, %	67.5	69.2	40.3	50.0
cORR, %	52.5	61.5	30.6	38.5
DCR, %	87.5	92.3	87.1	80.8
mDoR, mo	8.5	12.3	NR	NR
mPFS, mo	5.6	15.0	5.4	6.7



2L, second-line; CNS, central nervous system; cORR, confirmed objective response rate; cPR, confirmed partial response; CT, computed tomography; DCR, disease control rate; EGFR, epidermal growth factor receptor; mDoR, median duration of response; mPFS, median progression-free survival; NR, not reached; NSCLC, non-small cell lung cancer; ORR, objective response rate; PBC, platinum-based chemotherapy; PD, progressive disease; PR, partial response; Q3W, once every 3 weeks; SD, stable disease.
 Zhang L, et al. ESMO 2023. Abstract 1316MO. Zhang L, et al. ASCO 2023. Abstract 3001.

BL-B01D1: Safety in All Tumor Types

Overall Safety Summary	All Q3W (N = 369)
Median follow-up (months)	3.9
TEAE, n (%)	363 (98)
• ≥ Grade 3	249 (67)
• ≥ Grade 4	125 (34)
• Serious	142 (38)
• Associated with death	17 (5)
• Associated with dc	12 (3)
• Associated with delay	102 (28)
• Associated with reduction	50 (14)
TRAE, n (%)	351 (95)
• ≥ Grade 3	226 (61)
• ≥ Grade 4	115 (31)
• Serious	108 (29)
• Associated with death*	8 (2)

- One grade 2 ILD was observed

TRAE ≥15%, %	All Q3W (N = 369)		2.5 mg/kg D1D8 Q3W (N = 278)		4.5 mg/kg D1 Q3W (N = 40)	
	Any	≥G3	Any	≥G3	Any	≥G3
Leukopenia	65	32	61	27	73	33
Anemia	64	24	64	22	73	25
Neutropenia	59	36	53	29	70	45
Thrombocytopenia	55	28	53	27	58	23
Nausea	36	<1	33	1	40	0
Asthenia	31	<1	28	1	33	0
Decreased appetite	29	<1	26	<1	38	0
Alopecia	25	0	21	0	43	0
Stomatitis	25	1	22	1	28	3
Vomiting	22	1	20	1	33	3
Diarrhea	17	<1	15	<1	30	0
Skin disorders	17	<1	14	<1	25	3
Hypokalemia	15	2	16	1	5	3
Hypoalbuminemia	13	0	15	0	5	0


► In regards to safety with BL-B01D1, similar to HER3-DXd different chemotherapy-related side effects were seen including cytopenia, nausea, vomiting, stomatitis, and alopecia. In regards to treatment-emergent adverse events that led to treatment discontinuation, that was quite low at 3%. And although pneumonitis or interstitial lung disease was rare, it was identified in one case and it was grade 2.



*Septic shock (n = 3); pneumonia (n = 2); respiratory failure, myelosuppression, gastrointestinal infection (n = 1 each).
G, grade; ILD, interstitial lung disease; Q3W, once every 3 weeks; TEAE, treatment-emergent adverse event; TRAE, treatment-related adverse event.
Zhang L, et al. ESMO 2023. Abstract 1316MO. Zhang L, et al. ASCO 2023. Abstract 3001.

SHR-A2009: Tumor Response in Advanced Solid Tumors

All Solid Tumors	All Doses (n = 36)
ORR, n (%)	9 (25.0)
DCR, n (%)	26 (72.2)
Median DoR, mo (range)	7.0 (2.8-8.5)
6-mo PFS, % (95% CI)	46.4 (27.0-63.8)
Patients With NSCLC	All Doses (n = 30)
ORR, n (%)	9 (30.0)
DCR, n (%)	23 (76.7)
Median DoR, mo (range)	7.0 (2.8-8.5)
6-mo PFS, % (95% CI)	49.8 (28.8-67.8)



- Among patients with NSCLC (n = 36), 94.4% had an EGFR mutation and all were resistant to EGFR-TKI, with 85.3% (29/34) previously treated with third-generation agents
- Grade ≥3 TRAEs: 13 (31.0%), leading to drug discontinuation in 3 (7.1%) patients
 - Interstitial lung disease occurred in 2 (4.8%) patients

► A different agent is SHR-A2009, which is a different HER3-directed antibody-drug conjugate and this also was looked at in patients with non-small cell lung cancer and the vast majority had EGFR-mutations and were previously treated with a third-generation EGFR TKI. Response rate to this drug was 25% in 36 patients treated. And so also looked quite promising and this drug is development is currently ongoing.




DCR, disease control rate; DoR, duration of response; EGFR, epidermal growth factor receptor; NSCLC, non-small cell lung cancer; ORR, objective response rate; PFS, progression-free survival; TKI, tyrosine-kinase inhibitor; TRAE, treatment-related adverse event.
Zhou Q, et al. ESMO 2023. Abstract 698MO.

SHR-A2009: AE Profile

- No dose-limiting toxicities occurred up to 10.5 mg/kg Q3W dose level

Event, n (%)	All patients (n = 42)
Median duration of treatment, mo (range)	2.8 (0.3-12.4)
Any AE	42 (100)
Grade ≥3 AE	21 (50.0)
Any TRAE	39 (92.9)
Grade ≥3 TRAE	13 (31.0)
TRAE leading to dose reduction	3 (7.1)
TRAE leading to dose hold	8 (19.0)
TRAE leading to discontinuation	3 (7.1)
TRAE leading to death	1 (2.4)
Serious TRAE	4 (9.5)
ILD	2 (4.8)



AE, adverse event; ALP, alkaline phosphatase; ALT, alanine aminotransferase; AST, aspartate aminotransferase; GGT, gamma-glutamyl transferase; ILD, interstitial lung disease; Q3W, once every 3 weeks; TRAE, treatment-related adverse event; WB, white blood cell.
Zhou Q, et al. ESMO 2023. Abstract 658MO.


▶ In regards to adverse events, in this initial phase 1 there weren't any dose-limiting toxicities that were seen up to the highest dose assessed. In regards to treatment-related adverse events very similar profile with some cytopenia seen as well as nausea, vomiting, alopecia, and fatigue. And the rate of treatment-related adverse events leading to treatment discontinuation was 7.1%.

So, a lot of us have a lot of excitement of how these HER3-directed antibody-

drug conjugates will improve clinical outcomes for our patients with EGFR-mutant lung cancer. So, our standard of care treatment algorithm presently is to proceed with a third-generation EGFR TKI that could be with cytotoxic chemotherapy or without. After progression on osimertinib, if chemotherapy wasn't used in the first-line setting, you can use chemotherapy in the second-line setting, but after that there aren't any targeted therapies that are approved in that later-line setting. And

so our options really are just cytotoxic chemotherapy to have different antibody-drug conjugates including HER3-DXd. And some of these other agents for use in this space is really an exciting prospect and we look forward to the results of the study that's assessing HER3-DXd compared to platinum-based chemotherapy. Because there is that question about whether we can utilize these therapies prior to standard cytotoxic chemotherapies.

ADC Mechanisms of Toxicity



Cytotoxic payload:

- Intrinsic toxicity
- Drug-to-antibody ratio
- Membrane permeability (bystander effect)

Linker:

- Cleavable vs noncleavable

Tumor Antigen Target:

- Expression of target protein on noncancer cells

AXIS
Medical Education

ADC, antibody-drug conjugate.
Johns AC, Campbell MT. *Cancer J.* 2022;28(6):469-478.


► And so, when we think about treatment-emergent adverse events I think something that's critical, is to understand what the different class adverse events are and to be able to detect them properly and be able to manage those adverse events well.

And so, when we think about antibody-drug conjugates and mechanisms of toxicity, the primary toxicity is largely from the cytotoxic chemotherapy payload. That affects both the cell of interest, which is of course the tumor cell, but also can lead to bystander effect and release of that

chemotherapy payload to other neighboring cells. And then sometimes there can also be toxicities that are related to the antibody and so, for example, for HER2-directed antibodies you can see cardiotoxicity because of HER2 expression on cardiomyocytes.

ADC Mechanisms of Toxicity: Bystander Effect

- Internalized ADC degraded by target cell and payload exits cell and enters neighbor cell
- Payload released in extracellular space without ADC entering target cell (acidic)



AXIS
Medical Education

ADC, antibody-drug conjugate; ADCC, antibody-dependent cellular cytotoxicity; ADCP, antibody-dependent cellular phagocytosis. Johns AC, Campbell MT. *Cancer J.* 2022;28(6):469-478. Drago JZ, et al. *Nat Rev Clin Oncol.* 2021;18(6):327-344.

► Here is an illustration of the bystander effect where typically we see an ADC get internalized after binding to the target antigen and then should be entering the tumor cell and then cleaved with the payload being released while inside the tumor cell, but then they're can be payload released from

either the dying tumor cells or prematurely released that does kill adjacent cells and that's where we see some of the toxicity with these ADCs.

In regards to a very important toxicity that is, as I mentioned before, pneumonitis or interstitial lung disease, again

this is relatively rare but also a class effect of these deruxtecan backbone ADCs and it's just really critical to have a high suspicion for this, and to diagnose it early because patient outcomes are best when treatment is immediately implemented.

Workup for Suspected ADC-Related ILD

- Hold ADC pending more information
- History and physical exam
- Rule out other causes of ILD (eg, other drugs or RT toxicity) and other pathologies with similar presentation (eg, infection, PD, or PE)
 - High-resolution CT scan of chest
 - Pulmonology consult with pulmonary function testing
 - Bronchoscopy and BAL ± transbronchial lung biopsy
- Laboratory tests
 - CBC, liver and kidney function tests, electrolytes, CRP, ESR, procalcitonin, LDH, other
 - Analysis for infection based on suspected pathogen (blood culture, expectorated sputum, urinary antigens, β -D-glucan, other)
 - Tumor markers and autoimmune antibodies, if indicated



ADC, antibody-drug conjugate; BAL, bronchoalveolar lavage; CBC, complete blood count; CRP, C-reactive protein; CT, computed tomography; ESR, erythrocyte sedimentation rate; ILD, interstitial lung disease; LDH, lactate dehydrogenase; PD, progressive disease; PE, pulmonary embolism. Tarantino P, et al. *JAMA Oncol*. 2021;7(12):1873-1881.

► And so, in terms of workup for suspected ADC-related interstitial lung disease, I think the first thing is that, especially in lung cancer, it can be a very challenging diagnosis because the majority of our patients do have shortness of breath and cough and respiratory symptoms. But really, I counsel my patients if there is a

change from baseline in terms of your respiratory symptoms, I would just, be very quick to hold the antibody-drug conjugate while we're trying to gather more information. So, if you ever have a suspicion for ILD, you hold the antibody-drug conjugate. You look out for other causes of potential infectious or inflammatory

changes and that typically means potentially starting steroids or treating infection if appropriate. And then always referral to some of my colleagues like pulmonary to get a diagnostic biopsy or bronchoscopy can be really critical and, of course, further imaging to assess any radiographic findings.

Detecting and Managing T-DXd–Related Interstitial Lung Disease: The 5 “S” Rules



► And so, again, when we think about managing interstitial lung disease or pneumonitis it's screening, having an actual suspicion for ILD, scanning at first sign of change in symptoms, synergy with care teams, so reaching out to pulmonary or other colleagues, stopping treatment, and then steroids as our main stay of treatment for pneumonitis.



T-DXd, trastuzumab deruxtecan.
Taranfiro P, Tolanev SM. *JCO Oncol Pract.* 2023;19(8):526-527.

Management of ILD Associated With HER3-DXd




What to Look for

- Shortness of breath, particularly on exertion
- Dry cough
- Chest discomfort
- Fatigue



Promptly investigate any evidence of suspected ILD/pneumonitis with high-resolution CT, pulmonologist consult, and blood cultures/CBC



Grade and Description	Protocol Management Recommendations
1: asymptomatic; clinical or diagnostic observations only	<ul style="list-style-type: none"> • Hold patritumab deruxtecan until resolution to grade 0 <ul style="list-style-type: none"> • If AE resolves in ≤28 days, resume with same dose of patritumab deruxtecan • If AE resolves in >28 days, resume with reduced dose of patritumab deruxtecan • Consider corticosteroid treatment (eg, prednisone ≥0.5 mg/kg/day)
2: symptomatic; limiting instrumental ADL 3: severe symptoms; limiting self-care ADL or life-threatening respiratory compromise	<ul style="list-style-type: none"> • Permanently discontinue patritumab deruxtecan • Promptly initiate corticosteroid treatment (eg, ≥1 mg/kg/day prednisolone or equivalent) and continue for ≥14 days followed by gradual taper for ≥4 weeks
For all grades	<ul style="list-style-type: none"> • Oxygen supplementation for hypoxia • Monitor closely for worsening symptoms, re-image as clinically indicated • Supportive treatment for prolonged corticosteroid use • Consider infliximab, mycophenolate mofetil, IVIG, etc if corticosteroid refractory

Dose reductions: Starting Dose 5.6 mg/kg Q3W → Reduction 1: 4.8 mg/kg → Reduction 2: 3.2 mg/kg → Discontinue



Image courtesy of Rebecca Heist, MD, MPH.
ADL, activities of daily living; AE, adverse event; CBC, complete blood count; CT, computed tomography; HER3-DXd, patritumab deruxtecan;
ILD, interstitial lung disease.
Yu HA, et al. *J Clin Oncol.* 2023;41(35):5363-5375.
Janne PA, et al. *Cancer Discov.* 2022;12(7):74-89.

► And, typical and similar to grading of other adverse events, we can grade interstitial lung disease or pneumonitis by the CTCAE criteria. And so, if patients have asymptomatic pneumonitis where we see something on imaging when patients feel completely fine, we still hold patritumab deruxtecan, but once those radiographic findings resolve, it can be appropriate to resume patritumab deruxtecan or HER3-DXd. But importantly, if somebody has symptomatic interstitial lung disease or pneumonitis, we still have to do the same things; stop drug, initiate corticosteroids, think about treating other causes like infection but we don't tend to restart. And so the guidelines currently are if patients have grade 2 or greater pneumonitis, we permanently discontinue HER3-DXd.

Managing Clinically Significant Nausea and Vomiting With HER3-DXd

- Premedicate with 3-drug regimen for CINV (eg, dexamethasone + 5-HT₃ receptor antagonist + NK1 receptor antagonist)
- **Onset may be delayed:** Provide patient with take-home antiemetics (eg, dexamethasone, ondansetron)
- Manage with antiemetics, dose reductions; withhold if high grade until resolved to grade ≤1

AXIS
Medical Education

CINV, chemotherapy-induced nausea and vomiting; HER3-DXd, patritumab deruxtecan.
Stankovicz M, et al. *Breast Care (Basel)*. 2021;16(4):408-411.
Rugo HS, et al. *ESMO Open*. 2022;7(4):100553.
Trastuzumab deruxtecan. Package insert. Daiichi Sankyo, Inc; 2024.

► In terms of nausea and vomiting with HER3-DXd all of us are very familiar with this toxicity. We give platinum-based chemotherapy and other regimens that induce nausea and so it's really just using the appropriate medications that we use with our other emetogenic cancer therapies. So, dexamethasone, a 5-HT₃ receptor antagonist, and an NK-1 receptor antagonist. So and often times with these different ADCs I also do a prolonged steroid taper, which I think can be helpful, but definitely giving antiemetics with therapy is key to managing nausea.

Management of Select AEs Associated With HER3-DXd: Neutropenia

Grade/Description	Protocol Management Recommendations
Grade 1: <LLN - 1500 neutrophils/mm ³ ; <LLN - 1.5 x 10 ⁹ neutrophils/L	• Continue patritumab deruxtecan and monitor for worsen neutropenia
Grade 2: <1500 - 1000 neutrophils/mm ³ ; <1.5 - 1.0 x 10 ⁹ neutrophils/L	• Continue patritumab deruxtecan and monitor for worsen neutropenia
Grade 3: 500 to <1000 neutrophils/mm ³ ; 0.5-1 x 10 ⁹ neutrophils/L	• Hold patritumab deruxtecan until resolution to grade ≤2 • Then resume with same dose of patritumab deruxtecan
Grade 4: <500 neutrophils/mm ³ ; <0.5 x 10 ⁹ neutrophils/L	• Hold patritumab deruxtecan until resolution to grade ≤2 • Then resume with reduced dose of patritumab deruxtecan

Dose reductions: Starting Dose 5.6 mg/kg Q3W → Reduction 1: 4.8 mg/kg → Reduction 2: 3.2 mg/kg → Discontinue

AXIS
Medical Education

AE, adverse event; HER3-DXd, patritumab deruxtecan; Q3W, once every 3 weeks.
Yu HA, et al. *J Clin Oncol*. 2023;41(35):5363-5375.
Clinical Study Protocol. https://ascopubs.org/doi/suppl/10.1200/JCO.23.01476/suppl_file/protocol_JCO.23.01476.pdf

► In regards to neutropenia, again we're comfortable with this because we do give cytotoxic chemotherapy. Always worth it to hold or delay dosing if neutropenia is seen. If there's concern for infection, you can give growth factor support as intervention. And then if neutropenia is severe you can always think about giving prophylactic dose growth factor support like Neulasta with therapy.

Management of Select AEs Associated With HER3-DXd: Febrile Neutropenia

Grade/Description	Protocol Management Recommendations
Grade 3: ANC <1000/mm ³ with a single temperature of >38.3°C (101°F) or sustained temperature of ≥38°C (100.4°F) for ≥1 hr	<ul style="list-style-type: none"> Hold patritumab deruxtecan until resolution Then resume with patritumab deruxtecan and consider dose reduction Consider administration of G-CSF as prophylaxis for all subsequent cycles and according to local guidelines
Grade 4: Life-threatening consequences; urgent intervention indicated	<ul style="list-style-type: none"> Hold patritumab deruxtecan until resolution Then resume with reduced dose of patritumab deruxtecan Administer of G-CSF as prophylaxis for all subsequent cycles and according to local guidelines

Dose reductions: Starting Dose 5.6 mg/kg Q3W → Reduction 1: 4.8 mg/kg → Reduction 2: 3.2 mg/kg → Discontinue



AE, adverse event; ANC, absolute neutrophil count; HER3-DXd, patritumab deruxtecan; G-CSF, granulocyte colony-stimulating factor.
Yu HA, et al. J Clin Oncol. 2023;41(35):5363-5375.
Clinical Study Protocol. https://ascopubs.org/doi/suppl/10.1200/JCO.23.01476/suppl_file/protocol_JCO.23.01476.pdf

- ▶ Febrile neutropenia of course a more serious sequelae where we do have to give G-CSF and of course if there's any febrile neutropenia recommendation would be to administer G-CSF with further doses of HER3-DXd.

Management of Select AEs Associated With HER3-DXd: Thrombocytopenia

Grade/Description	Protocol Management Recommendations
Grade 1: <LLN - 75,000 platelets/mm ³ ; <LLN - 75.0 x 10 ⁹ platelets/L	<ul style="list-style-type: none"> Continue patritumab deruxtecan and monitor for worsen neutropenia
Grade 2: <75,000 - 50,000 platelets/mm ³ ; <75.0 - 50.0 x 10 ⁹ platelets/L	<ul style="list-style-type: none"> Continue patritumab deruxtecan and monitor for worsen neutropenia
Grade 3: <50,000 - 25,000/mm ³ platelets; <50.0 - 25.0 x 10 ⁹ platelets/L	<ul style="list-style-type: none"> Hold patritumab deruxtecan until resolution to grade ≤1 <ul style="list-style-type: none"> If AE resolves in ≤14 days, resume with same dose If AE resolves in >14 days, resume but consider reduced dose
Grade 4: <25,000 platelets/mm ³ ; <25.0 x 10 ⁹ platelets/L	<ul style="list-style-type: none"> Hold patritumab deruxtecan until resolution to grade ≤1 Then resume with reduced dose of patritumab deruxtecan

Dose reductions: Starting Dose 5.6 mg/kg Q3W → Reduction 1: 4.8 mg/kg → Reduction 2: 3.2 mg/kg → Discontinue



AE, adverse event; ANC, absolute neutrophil count; HER3-DXd, patritumab deruxtecan; LLN, lower limit of normal.
Yu HA, et al. J Clin Oncol. 2023;41(35):5363-5375.
Clinical Study Protocol. https://ascopubs.org/doi/suppl/10.1200/JCO.23.01476/suppl_file/protocol_JCO.23.01476.pdf

- ▶ And thrombocytopenia, again it typically is not associated with clinical sequelae, and you would just need to hold and restart when platelets are at a normal level.

So when I think about the toxicity or the safety profile of HER3-DXd it really is just being aware of symptoms and having a very low threshold for imaging because of that concern for pneumonitis or interstitial lung disease and then just managing the nausea, the fatigue, like we do with chemotherapy supportive measures to help insure that our patients have the best experience on these novel therapies.

The Importance of the MDT in NSCLC Care

- The wide range of treatment modalities requires collaboration among multiple specialists to develop individualized management strategies and provide optimal staging in the complex NSCLC setting
- The lung cancer MDT includes a medical oncologist, thoracic surgeon, pulmonologist, and radiation oncologist
 - Other specialists (eg, radiologists, pathologists, nurse navigator, nutritionists, nuclear medicine specialists, clinical pharmacists, molecular biologists, psychologists) may also be included
- **MDT-based patient care in NSCLC has been associated with longer overall survival and better quality-of-care-related outcomes**



NSCLC, non-small cell lung cancer; MDT, multi-disciplinary team.
De Castro Jr G, et al; for the Grupo Brasileiro de Oncologia Toracica. *JTO Clin Res Rep.* 2023;4(12):1-20.

► And so finally, thinking about shared decision-making and individualized treatment planning, as we all know this is really critical for a lung cancer and for a, when patients have a terminal diagnosis, understanding what their goals are with treatment is really critical. And so, there are, as we know, a wide range of treatment modalities and

options and so it really is key to have both buy-in from the patient as well as multiple specialists in order to provide individualized treatment plans for our patients. And that might be considering involvement of a thoracic surgeon, say if we need a PleurX placed or pleurodesis, radiation oncologist if there is a painful bone metastases,

and so all of these different specialists can be really critical as part of our treatment plan for our patients. And we know, that given this multi-disciplinary treatment, is associated with longer overall survival as well as better quality of life outcomes for our patients and so, really critical.

Treatment Shared Decision-Making (SDM) in NSCLC

In recent years, treatment options for NSCLC have rapidly expanded to include novel immunotherapies, targeted therapies, and combination and multidisciplinary approaches

Patients and their families are bombarded by multitudes of information about cancer, especially through social media and the internet

The complexity of cancer care and the abundance of cancer-related information can complicate the development of individualized care plans



NSCLC, non-small cell lung cancer.
Shickh S, et al. *Am Soc Clin Oncol Educ Book*. 2023;43:e389516.

▶ And I think this is a particularly relevant in EGFR-mutant lung cancer where there are oral therapies as well as intravenous therapies, understanding what people's goals are, how able they are to come to the clinic for intravenous treatments. Whether they are concerned about alopecia because of ongoing work commitments really becomes critical. And as the complexity of cancer care increases it's going to be even more important when our patients have more options to really understand what they value and to shape our treatment plans based on their values.

The Role of SDM in Planning Treatment Regimens

- SDM is a fundamental method of care that is central to individualizing treatment
- It involves an MDT approach to ensure optimal care for and communication with the patient and their family
- The initial step involves promoting productive dialogues that encourage active patient-clinician collaboration, facilitating the process of care plan development, and supporting the cocreation of a comprehensive care plan



MDT, multi-disciplinary team; SDM, shared decision-making.
Shickh S, et al. *Am Soc Clin Oncol Educ Book*. 2023;43:e389516.

▶ And so again, shared decision-making is a fundamental method of care that's central to our individualization of treatment. It ensures optimal care and communication with our patients' and is really, just something that we practice and do on a day-to-day basis. That really is important in our relationships with our patients. And so again, why do I think that this is important? I think it's important because it allows us to tailor our treatment to the goals and needs of individual patients because an 88-year-old patient that lives in a nursing home is very different than a 36-year-old with young children, who's married that, we're managing with the same disease, say EGFR-mutant lung cancer, and so it's pretty intuitive to state that we won't treat those patients the same and so we do the shared decision-making oftentimes even without acknowledging it overtly.

Case-Based Learning Lab

AXIS
Medical Education

- ▶ So to end with a few case studies.

Case Study 1

- 46-year-old woman with EGFR exon 19 deletion positive lung cancer who was initially started on osimertinib plus chemotherapy and had an initial good response but subsequent progression 22 months later
- She has multi-site progression in the liver, bone, and lung
- Repeat biopsy shows continued EGFR exon 19 deletion but no additional acquired genomic alterations

- ▶ Case study 1 is a 46-year-old woman with EGFR Exon 19 deletion-positive lung cancer who initially was started on osimertinib with chemotherapy, based on the FLAURA2 regimen, had an initial good response, but subsequently had progression. And this was multisite progression in the liver, bone, and lung and a repeat biopsy showed no additional acquired mutations.

AXIS
Medical Education

EGFR, epidermal growth factor receptor.

Case Study Question

What is the next best treatment option for this patient?

- a) Single agent docetaxel
- b) Carboplatin, pemetrexed, and amivantamab
- c) Osimertinib + capmatinib
- d) Patritumab deruxtecan



► So, what is the next best treatment option for this patient? I think this question has several options that could be potentially correct. Single agent docetaxel certainly is a second-line chemotherapy. I probably wouldn't choose carboplatin-pemetrexed or amivantamab because of the recent use of carboplatin and pemetrexed in the first-line setting. The question notes that there was no acquired mutation, so no acquired MET amplification, so I would not use capmatinib.

Case Study Question

What is the next best treatment option for this patient?

- a) Single agent docetaxel
- b) Carboplatin, pemetrexed, and amivantamab
- c) Osimertinib + capmatinib
- d) Patritumab deruxtecan



► But in the setting of prior osimertinib and prior platinum-based chemotherapy, if approved, my preferred choice would likely be patritumab deruxtecan and the idea is that the approval will be in this setting.

Case Study 2

- 63-year-old woman with L858R positive lung cancer who has metastases to bone, brain, and lung
- She was initially treated with osimertinib for 15 months, followed by carboplatin/pemetrexed for 8 months, and then develops progression in the lung with a new pleural effusion, dyspnea on exertion and a dry cough
- She is started on patritumab deruxtecan and after 3 cycles, her dyspnea on exertion and cough resolved
- Imaging shows resolution of the pleural effusion and shrinkage of her pulmonary mets
- She presents for cycle 6 of HER3-DXd with new onset dyspnea and with oxygen saturation of 87% on room air. She admits to a productive cough with some yellow sputum

AXIS
Medical Education

HER3-DXd, patritumab deruxtecan.

▶ The second case study was a 63-year-old woman with L858R-positive EGFR-mutant lung cancer with metastases to bone, brain, and lung. She was treated with osimertinib, but then developed clinical progression and then was treated with carboplatin and pemetrexed. She then developed a new pleural effusion as well as dyspnea upon exertion and then a dry cough. She was started in the third-line setting on patritumab deruxtecan and after 3 cycles had a marked improvement in her symptoms with improvement of dyspnea, her cough resolved, imaging showed resolution of her pleural effusion, as well as shrinkage of her pulmonary metastases. And then when she presented for cycle 6 of HER3-DXd she had new onset dyspnea over the last week, and when you checked oxygen saturation it was 87% on room air. She also had a productive cough with some yellow sputum.

Case Study Audience Question

What is the appropriate next step for this patient?

- a) Hold patritumab deruxtecan
- b) Start antibiotics for possible pneumonia
- c) Start prednisone at 1mg/kg
- d) Refer to pulmonary for workup and evaluation
- e) All of the above

▶ What is the appropriate next step for this patient? In this case the correct answer is all of the above. Any time there is a concern for ILD or pneumonitis I would immediately hold the potential offending agent, patritumab deruxtecan. You want to treat for other potential causes like infection, so starting antibiotics. Low threshold to start steroids in the setting of these symptoms. And then I would want to refer to pulmonary for further workup and consideration for bronchoscopy or further intervention.

AXIS
Medical Education

Key Takeaways

- HER3-directed ADC therapies may be a new treatment option for EGFR-mutated NSCLC with prior EGFR TKI exposure
 - ORR was similar for patients regardless of the type of prior EGFR TKI
- Management of TEAEs related to HER3-directed ADCs requires a multidisciplinary approach, with proactive monitoring for ILD/pneumonitis and early management of nausea/vomiting
- SDM is crucial to ensure that the patient's goal of therapy are included when selecting treatment



ADC, antibody-drug conjugate; EGFR, epidermal growth factor receptor; ILD, interstitial lung disease; SDM, shared decision-making; TEAE, treatment-emergent adverse event.

▶ The key takeaways are that HER3-directed antibody-drug conjugate therapies may be a new treatment option for EGFR-mutant lung cancer with prior EGFR TKI exposure and importantly the response, or efficacy, was similar with different mechanisms or resistance to

targeted therapy. Management of treatment-emergent adverse events really requires a multi-disciplinary approach with proactive monitoring for pneumonitis and early management of other symptoms like nausea and vomiting. And then finally, shared decision-making is, of

course, crucial to ensure that our patient's goals of therapy are included when selecting treatment.

Thanks very much. With that, we will conclude today's activity and thank you so much for your participation.

REFERENCES

- Daiichi Sankyo, Inc. Patritumab deruxtecan BLA submission receives complete response letter from FDA due to inspection findings at third-party manufacturer. Press release. https://www.daiichisankyo.com/files/news/pressrelease/pdf/202406/20240626_E.pdf
- Daiichi Sankyo, Inc. Patritumab deruxtecan granted U.S. FDA breakthrough therapy designation in patients with metastatic EGFR-mutated non-small cell lung cancer. Press release. https://www.daiichisankyo.com/files/news/pressrelease/pdf/202112/20211223_E1.pdf
- De Castro Jr G, Souza FH, Lima J, et al; for the Grupo Brasileiro de Oncologia Toracica. Does multidisciplinary team management improve clinical outcomes in NSCLC? A systematic review with meta-analysis. *JTO Clin Res Rep*. 2023;4(12):1-20.
- Dokala A, Thakur SS. Extracellular region of epidermal growth factor receptor: a potential target for anti-EGFR drug discovery. *Oncogene*. 2017;36(17):2337-2344.
- Drago JZ, Modi S, Chandarlapaty S. Unlocking the potential of antibody-drug conjugates for cancer therapy. *Nat Rev Clin Oncol*. 2021;18(6):327-344.
- Haratani K, Yonesaka K, Takamura S, et al. U3-1402 sensitizes HER3-expressing tumors to PD-1 blockade by immune activation. *J Clin Invest*. 2020;130(1):374-388.
- Hashimoto Y, Koyama K, Kamai Y, et al. A Novel HER3-Targeting Antibody-Drug Conjugate, U3-1402, Exhibits Potent Therapeutic Efficacy through the Delivery of Cytotoxic Payload by Efficient Internalization. *Clin Cancer Res*. 2019;25(23):7151-7161.
- HERTHENA-Lung01: A Phase 2 Randomized Open-Label Study of Patritumab Deruxtecan (U3-1402) in Subjects with Previously Treated Metastatic or Locally Advanced EGFR-mutated Non-Small Cell Lung Cancer (NSCLC) (Patritumab Deruxtecan in Subjects with Metastatic or Locally Advanced EGFR-mutated NSCLC). Clinical Study Protocol. Protocol Number: U31402-A-U201; IND Number 133343; EudraCT Number 2020-000730-17. Version 3.0, March 31, 2021. https://ascopubs.org/doi/suppl/10.1200/JCO.23.01476/suppl_file/protocol_JCO.23.01476.pdf
- HERTHENA-Lung02: a study of patritumab deruxtecan versus platinum-based chemotherapy in metastatic or locally advanced EGFRm NSCLC after failure of EGFR TKI therapy. ClinicalTrials.gov identifier: NCT05338970. Updated April 19, 2024. <https://clinicaltrials.gov/study/NCT05338970>
- Jänne PA, Baik C, Su W-C, et al. Efficacy and safety of patritumab deruxtecan (HER3-DXd) in EGFR inhibitor-resistant, EGFR-mutated non-small cell lung cancer. *Cancer Discov*. 2022;12(7):74-89.
- Johns AC, Campbell MT. Toxicities from antibody-drug conjugates. *Cancer J*. 2022;28(6):469-478.
- Jordan EJ, Kim HR, Arcila ME, et al. Prospective comprehensive molecular characterization of lung adenocarcinomas for efficient patient matching to approved and emerging therapies. *Cancer Discov*. 2017;7(6):596-609.
- Koganemaru S, Kuboki Y, Koga Y, et al. U3-1402, a novel HER3-targeting antibody-drug conjugate, for the treatment of colorectal cancer. *Mol Cancer Ther*. 2019;18(11):2043-2050.
- Kumagai X, Shao Y, Qin H-F, Tai Y-H, Gao H-J. ALK-rearrangement in non-small-cell lung cancer (NSCLC). *Thorac Cancer*. 2018;9(4):423-430.
- Lyu H, Han A, Polsdofer E, Liu S, Liu B. Understanding the biology of HER3 receptor as a therapeutic target in human cancer. *Acta Pharm Sin B*. 2018;8(4):503-510.
- Mok TSK, Jänne PA, Nishio M, et al. HERTHENA-Lung02: phase III study of patritumab deruxtecan in advanced EGFR-mutated NSCLC after a third-generation EGFR TKI. *Future Oncol*. 2024;20(15):969-908.
- Nakada T, Sugihara K, Jikoh T, Abe Y, Agatsuma T. The latest research and development into the antibody-drug conjugate, [fam-] trastuzumab deruxtecan (DS-8201a), for HER2 cancer therapy. *Chem Pharm Bull (Tokyo)*. 2019;67(3):173-185.
- Ogitani Y, Aida T, Hagihara K, et al. DS-8201a, A novel HER2-targeting ADC with a novel DNA topoisomerase I inhibitor, demonstrates a promising antitumor efficacy with differentiation from T-DMI. *Clin Cancer Res*. 2016;22(20):5097-5108.
- Ogitani Y, Hagihara K, Oitate M, Naito H, Agatsuma T. Bystander killing effect of DS-8201a, a novel anti-human epidermal growth factor receptor 2 antibody-drug conjugate, in tumors with human epidermal growth factor receptor 2 heterogeneity. *Cancer Sci*. 2016;107(7):1039-1046.
- Ramalingam SS, Gray JE, Ohe Y, et al. Osimertinib vs comparator EGFR-TKI as first-line treatment for EGFRm advanced NSCLC (FLAURA): Final overall survival analysis. Abstract presented at: European Society for Medical Oncology Annual Meeting; Barcelona, Spain; September 27-October 1, 2019. Abstract 567.
- Rugo HS, Bianchini G, Cortes J, Henning J-W, Untch M. Optimizing treatment management of trastuzumab deruxtecan in clinical practice of breast cancer. *ESMO Open*. 2022;7(4):100553.
- Scharpenseel H, Hanssen A, Loges S, et al. EGFR and HER3 expression in circulating tumor cells and tumor tissue from non-small cell lung cancer patients. *Sci Rep*. 2019;9:7406.
- Shickh S, Leventakos K, Lewis MA, Bombard Y, Montori VM. Shared decision making in the care of patients with cancer. *Am Soc Clin Oncol Educ Book*. 2023;43:e389516.
- Sithanandam G, Anderson LM. The ERBB3 receptor in cancer and cancer gene therapy. *Cancer Gene Ther*. 2008;15(7):413-448.
- Soria J-C, Ohe Y, Vansteenkiste J, et al; for the FLAURA Investigators. Osimertinib in untreated EGFR-mutated advanced non-small-cell lung cancer. *N Engl J Med*. 2018;378:113-125.
- Stankowicz M, Mauro L, Harnden K, Pennisi A. Management of chemotherapy-induced nausea and vomiting with trastuzumab deruxtecan: a case series. *Breast Care (Basel)*. 2021;16(4):408-411.
- Tarantino P, Modi S, Tolaney S, et al. Interstitial lung disease induced by anti-ERBB2 antibody-drug conjugates: A review. *JAMA Oncol*. 2021;7(12):1873-1881.
- Tarantino P, Tolaney SM. Detecting and managing T-DXd-related interstitial lung disease: the five "S" rules. *JCO Oncol Pract*. 2023;19(8):526-527.
- Trastuzumab deruxtecan. Package insert. Daiichi Sankyo, Inc; 2024.
- Yu H, Goto Y, Hayashi H, et al. Patritumab deruxtecan (HER3-DXd) in EGFR-mutated NSCLC following EGFR TKI and platinum-based chemotherapy: HERTHENA-Lung01. Abstract presented at: IASLC World Conference on Lung Cancer; Singapore; September 9-12, 2023. Abstract OA05.03.

REFERENCES

- Yu HA, Goto Y, Hayashi H, et al. HERTHENA-Lung01, a phase ii trial of patritumab deruxtecan (HER3-DXd) in epidermal growth factor receptor-mutated non-small-cell lung cancer after epidermal growth factor receptor tyrosine kinase inhibitor therapy and platinum-based chemotherapy. *J Clin Oncol*. 2023;41(35):5363-5375.
- Zhang L, Ma Y, Zhao Y, et al. BL-B01D1, a first-in-class EGFRxHER3 bispecific antibody-drug conjugate, in patients with non-small cell lung cancer: Updated results from first-in-human phase I study. Abstract presented at: European Society for Medical Oncology Annual Meeting; Madrid, Spain; October 20-24, 2023. Abstract 1316MO.
- Zhang L, Ma Y, Zhao Y, et al. BL-B01D1, a first-in-class EGFRxHER3 bispecific antibody-drug conjugate (ADC), in patients with locally advanced or metastatic solid tumor: Results from a first-in-human phase 1 study. Abstract presented at: American Society of Clinical Oncology Annual Meeting; Chicago, Illinois; June 2-6, 2023. Abstract 3001.
- Zhou Q, Wu Y, Li J, et al. Phase I study of SHR-A2009, a HER3-targeted ADC, in advanced solid tumors. Abstract presented at: European Society for Medical Oncology Annual Meeting; Madrid, Spain; October 20-24, 2023. Abstract 658MO.

About AXIS Medical Education, Inc.

AXIS Medical Education, Inc. is a full-service continuing education company that designs and implements live, web-based, and print-based educational activities for healthcare professionals. AXIS provides convenient opportunities to engage learners based on their individual learning preferences through a full spectrum of educational offerings.

The executive leadership of AXIS combines 75 years of experience in adult learning theory, curriculum design/implementation/assessment, continuing education accreditation standards, and medical meeting planning and logistics. Our team has a deep understanding of the governing guidelines overseeing the medical education industry to ensure compliant delivery of all activities.

AXIS employs an experienced team of medical and scientific experts, medical writers, project managers, meeting planners, and logistics professionals. This team is dedicated to meeting the unmet educational needs of healthcare professionals, with the goal of improving patient outcomes.

AXIS believes that partnerships are crucial in our mission to deliver timely, relevant, and high-quality medical education to healthcare professionals. To that end, AXIS partners with other organizations and accredited providers to offer added expertise and assist in expanding access to our educational interventions. AXIS also partners with numerous patient advocacy organizations to provide recommended patient education and caregiver resources in specific disease areas. AXIS finds value in these partnerships because they complement our core clinical curriculum with validated and relevant supplemental resources for busy clinicians and their patients.

The mission of AXIS is to enhance the knowledge, skills, competence, and performance of the interprofessional healthcare team to ensure patients receive quality care, resulting in improved patient outcomes. We engage healthcare professionals in fair-balanced, scientifically rigorous, expert-led certified educational activities designed to foster lifelong learning that is applicable to clinical practice and patient-centered care.

To learn more and to see our current educational offerings, visit us online at www.AXISMedEd.com.

